


Practice Paper 1 (Pure 1)

Scan here to return to the course
or visit [savemyexams.com](https://www.savemyexams.com)



Total Marks

/75

- 1 (a)** The fifth term of an arithmetic series is k , where k is a constant, and the sum of the first eight terms of the series is $2k$.

Show that the first term, a , of the series is $-5k$.

(3 marks)

- (b)** Find an expression for the common difference, d , of the series in terms of k .

(2 marks)

2 (a) Find the first three terms in the expansion of $(4 - 3x)^9$.

(3 marks)

(b) Given that x is small such that x^3 and higher powers of x can be ignored show that

$$(3 - 2x^2)(4 - 3x)^9 \approx 786432 - 5308416x + 15400960x^2$$

(3 marks)

- 3 (a)** A ball is dropped and bounces such that the height of each bounce is 80% of the height of the previous bounce.

The first bounce of the ball reaches a height of 1.60 m.

Find the height the ball bounces on its 8th bounce.

(2 marks)

- (b)** Find the total vertical distance travelled by the ball from when it first hits the ground until it hits the ground at the end of its twentieth bounce.

(2 marks)

- (c)** Give one reason why this model may be unrealistic.

(1 mark)

- 4 (a)** The point $P(3, -12)$ lies on the curve with equation $y = x^2 - 12x + 15$.

The graph is stretched so that the point P is mapped to the point $(3, -4)$. Write down the equation of the transformed function in the form $y = ax^2 + bx + c$, where a , b and c are constants to be found.

(2 marks)

- (b)** The graph is stretched so that the point P is mapped to the point $(1, -12)$. Write down the equation of the transformed function in the form $y = (dx)^2 - 12(dx) + 15$, where d is a constant to be found.

(2 marks)

5 (a) The curve C has equation $y = x^2 - 3x + 2$. The line l has equation $y = 3x - 7$.

Find any points of intersection between C and l .

(3 marks)

(b) Sketch the graphs of C and l , showing clearly any points of intersection with the coordinate axes for both graphs, the minimum point of C and any points of intersection found between C and l .

(3 marks)

6 (a) The functions $f(x)$ and $g(x)$ are defined as follows

$$f(x) = 3x^2 + 2 \quad x \in \mathbb{R}$$

$$g(x) = 1 - 3x \quad x \in \mathbb{R}$$

Write down the range of $f(x)$.

(1 mark)

(b) Find

(i) $fg(x)$


(ii) $gf(x)$

(4 marks)

(c) Solve the equation $f(x) = g(x) + 1$

(2 marks)

7 The graph below shows the curve with equation $y = 2 \sin \theta$, in the interval $0^\circ \leq \theta \leq 360^\circ$. One value of θ has been labelled ($\theta = a^\circ$).



Use the graph, along with the symmetry properties of the sine function, to verify that


$$2 \sin a = 2 \sin(180^\circ - a) = -2 \sin(180^\circ + a) = -2 \sin(360^\circ - a).$$

(2 marks)

8 Solve the equation $2 \sin 2\theta = 1$ for $0 \leq \theta \leq 2\pi$.

(3 marks)

9 (a) The diagram below shows the sector of a circle OAB .



Show that the area of the shaded segment is given by $\frac{1}{2}r^2(\theta - \sin \theta) \text{ cm}^2$.

(3 marks)

(b) Find, in terms of θ , the percentage of the sector that the segment occupies.

(2 marks)

10 (a) A curve is described by the equation $y = f(x)$, where $f(x) = 7 - 2x^2 + \sqrt{x}$, $x \geq 0$.

Find $f'(x)$ and $f''(x)$.

(3 marks)


(b) P is the stationary point on the curve.

Find the coordinates of P and determine its nature.

(4 marks)

11 A circle has equation $x^2 + y^2 + 14x - 6y = -41$.


The lines l_1 and l_2 are both tangents to the circle, and they intersect at the point $(0, 14)$.



Find the equations of l_1 and l_2 , giving your answers in the form $y = mx + c$.

(7 marks)

- 12 (a)** The line with equation $5y = 14 - 3x$ cuts the curve with equation $5y = 20 - 2x - x^2$ at the points P and Q , as shown.



Find the x - and y -coordinates of the points P and Q .

(5 marks)

- (b)** Find the exact area of the region labelled R , giving your answer in the form $\frac{a}{b}$, where a and b are integers to be found.

(6 marks)

- 13 (a)** A mathematical model for a bowl is obtained by rotating through 360° about the y -axis the part of the curve $y = \frac{1}{a}x^2$ which is between $y = 8$ and $y = 9$ and then adding a flat bottom. a is a constant such that $a > 0$.

Find the capacity of the bowl in terms of a .

(4 marks)

- (b)** In the case where the boundaries $y = 3$ and $y = 6$ are used in place of $y = 8$ and $y = 9$, the capacity of the bowl is increased by 15π cubic units. Find the value of a .

(3 marks)